






ORIGINAL RESEARCH

Growth, longevity and mortality of pink shrimps *Farfantepenaeus brasiliensis* and *F. paulensis* in southeastern Brazil

JÚLIA FERNANDES FERROCA^{1*}, LIZANDRA FERNANDES MIAZAKI¹, ADILSON FRANSOZO², FÚLVIO AURÉLIO DE MORAES FREIRE³, ROGERIO CAETANO COSTA¹

¹Laboratory of Biology of Marine and Freshwater Shrimps (LABCAM), Biological Sciences Department, Faculty of Sciences, São Paulo State University, Bauru, Brazil. ²Study Group on Crustacean Biology, Ecology and Culture (NEBECC) – Zoology Department, Bioscience Institute, UNESP, 18618-000, Botucatu, Brazil. ³Laboratory of Biology, Ecology and Evolution of Crustaceans (LABECC), Department of Botany and Zoology, UFRN, Lagoa Nova University Campus, 59078-970, Natal/RN – Brazil. ORCID *Julia Fernandes Ferroca*  <https://orcid.org/0000-0002-7274-3941>, *Lizandra Fernandes Miazaki*  <https://orcid.org/0000-0002-1678-3228>, *Adilson Fransozo*  <https://orcid.org/0000-0002-2067-5406>, *Fúlvio Aurélio De Moraes Freire*  <https://orcid.org/0000-0003-1580-0222>, *Rogério Caetano Da Costa*  <https://orcid.org/0000-0002-1342-7340>.



ABSTRACT. The study estimated the parameters of growth, longevity, fishing mortality (F), natural mortality (M) and total mortality (Z) and the exploitation rates (E) of the shrimps *Farfantepenaeus brasiliensis* and *F. paulensis* sampled in Ubatuba Bay from January to December 2000. Shrimps were identified by species, sexed and measured (carapace length - CL in mm). Overall, 1,231 individuals of *F. brasiliensis* and 687 of *F. paulensis* were analyzed. The mean size between sexes did not differ for both species. The estimated parameters of *F. brasiliensis* were: $CL_{\infty} = 41.08$ mm, $k = 2.41$ year⁻¹ for males and $CL_{\infty} = 47.32$ mm, $k = 2.23$ year⁻¹ for females; longevity of 1.91 years (males) and 2.05 years (females); M of 2.47 (males) and 2.28 (females); F of 7.97 (males) and 8.42 (females). For *F. paulensis*, the following values were observed: $CL_{\infty} = 36.55$ mm, $k = 2.41$ year⁻¹ for males and $CL_{\infty} = 49.24$ mm, $k = 2.51$ year⁻¹ for females; longevity of 1.91 years (males) and 1.81 years (females); M of 2.52 (males) and 2.52 (females); F of 7.64 (males) and 10.25 (females). The high values of k and F found for both species compared to those from the literature reflected the high E values, indicating that at the time, the closed season was still not responsible for stock recovery. We highlight the need for studies to assess the current status of stocks so they can be compared to the results found herein.

Keywords: Artisanal fishery, Bertalanffy, closed season, exploitation rate, Penaeidae.

Crecimiento, longevidad y mortalidad de camarones rosados *Farfantepenaeus brasiliensis* y *F. paulensis* en el sureste de Brasil

RESUMEN. El estudio estimó los parámetros de crecimiento, longevidad, mortalidad por pesca (F), mortalidad natural (M) y mortalidad total (Z) y las tasas de explotación (E) de los camarones *Farfantepenaeus brasiliensis* y *F. paulensis* muestreados en la Bahía de Ubatuba de enero a diciembre de 2000. Los camarones fueron identificados por especie, sexados y medidos (longitud del caparazón - CL en mm). En total, se analizaron 1.231 individuos de *F. brasiliensis* y 687 de *F. paulensis*. La talla media entre sexos no difirió para ambas especies. Los parámetros estimados de *F. brasiliensis* fueron: $CL_{\infty} = 41,08$ mm, $k = 2,41$ año⁻¹ para machos y $CL_{\infty} = 47,32$ mm, $k = 2,23$ año⁻¹ para hembras; longevidad de 1,91 años (machos) y 2,05 años (hembras); M de 2,47 (machos) y 2,28 (hembras); F de 7,97 (machos) y 8,42 (hembras). Para *F. paulensis* se observaron los siguientes valores: $CL_{\infty} = 36,55$ mm, $k = 2,41$ año⁻¹ para machos y $CL_{\infty} = 49,24$ mm, $k = 2,51$ año⁻¹ para hembras; longevidad de 1,91 años (machos) y 1,81 años (hembras); M de 2,52 (machos) y 2,52 (hembras); F de 7,64 (machos) y 10,25 (hembras). Los altos valores de k y F encontrados para ambas especies en comparación con los de la literatura reflejaron los altos valores de E, lo que indica que en ese momento la temporada de veda todavía no era responsable de la recuperación del stock. Resaltamos la necesidad de estudios que evalúen el estado actual de los stocks para que puedan ser comparados con los resultados aquí encontrados.



*Correspondence:
juliaperroca@gmail.com

Received: 28 October 2021
Accepted: 12 December 2021

ISSN 2683-7595 (print)
ISSN 2683-7951 (online)

<https://ojs.inidep.edu.ar>

Journal of the Instituto Nacional de
Investigación y Desarrollo Pesquero
(INIDEP)



This work is licensed under a Creative
Commons Attribution-
NonCommercial-ShareAlike 4.0
International License

Palabras clave: Pesquería artesanal, Bertalanffy, cierre de temporada, tasa de explotación, Penaeidae.

INTRODUCTION

Farfantepenaeus brasiliensis (Latreille, 1817) and *F. paulensis* (Pérez-Farfante, 1967), commonly known as pink shrimps, are highly exploited fishing resources on the southern and southeastern coast of Brazil (Costa and Fransozo 1999; Teodoro et al. 2016). Both species are exploited by two fleet types during different stages of their life cycles, i.e., industrial fleets offshore exploit adults in the open sea, while artisanal fisheries reach juveniles in estuarine environments and coastal lagoons (D’Incao 1991).

Accurate information about the age of the target organisms in fisheries is essential for calculating parameters such as longevity, growth rate and mortality, which end up reflecting the human impacts on natural populations (Kilada and Driscoll 2017). The most common method for studying age in natural populations is based on the analysis of distribution of length frequencies, in which the raw data is grouped into class intervals and, subsequently, algorithms are used to obtain the normal distribution peaks (e.g., in each month of collection) (Castilho et al. 2015; Silva et al. 2015). Finally, modal groups are detected and their combination supports the final model adjusted by the von Bertalanffy curve (1938). According to Vogt (2019), the von Bertalanffy curve makes it possible to estimate longevity from frequency data and life history.

Some growth studies have been conducted along the Brazilian coast for Penaeidae species, namely: *Artemesia longinaris* Spence Bate, 1888 (Semensato and Di Benedetto 2008), *F. brasiliensis* (Mello 1973; Vilela et al. 1997; Leite Jr and Petrere Jr 2006; Lopes 2012), *F. paulensis* (Mello 1973; D’Incao 1984; Leite Jr and Petrere Jr 2006; Lopes 2012), *Farfantepenaeus subtilis* (Pérez-Farfante, 1967) (Silva et al. 2015; Santos et al. 2020), *Litopenaeus schmitti* (Burkenroad, 1936) (Miazaki et al. 2018; Santos et al. 2020; Carvalho et al. 2021), and *Xiphopenaeus kroyeri* (Heller, 1862) (Branco et al. 1994; Grabowski et al. 2014; Lopes et al. 2014; Castilho et al. 2015; Silva et al. 2018).

Studies concerning pink-shrimps, *F. brasiliensis* and *F. paulensis*, were conducted either in Santos/Guarujá (central coast of São Paulo state) or in coastal lagoons in Rio de Janeiro State

(Araruama Lagoon) and Rio Grande do Sul State (Patos Lagoon). These areas differ significantly from our study area, Ubatuba Bay (northern coast of São Paulo state). Araruama Lagoon is the largest permanently hypersaline lagoon in the world (Souza et al. 2003), while Patos Lagoon is the largest coastal lagoon in the world (Kjerfve 1986). The Santos/Garujá area is located in an estuarine complex with extensive mangrove formations (Lamparelli et al. 2001). Ubatuba Bay, however, has small estuaries that are susceptible to changes in salinity forcing *Farfantepenaeus* spp. juveniles to use shallow portions of the bay as auxiliary nursing areas (Costa et al. 2008).

In addition to calculating growth parameters, it is necessary to calculate both natural and fishing mortality, monitor stocks and elaborate adequate management plans which will guarantee the long-term maintenance of these stocks (King 2007) and consider their economic, ecological and social importance (Musiello-Fernandes et al. 2017). In the southeastern region, both species are frequently targeted due to their large sizes and high commercial value. They are also captured as bycatch of the seabob shrimp *X. kroyeri* in shallow marine areas (Mantelatto et al. 2016), whose population in Ubatuba is already overexploited (Miazaki et al. 2021). We believe these factors may be influencing and shaping populational parameters of pink shrimps in the sampling area. Thus, the present study used a length frequency analysis to estimate the (i) growth and longevity, (ii) total, natural and fishing mortality, and (iii) the exploitation rates of both sexes of *F. brasiliensis* and *F. paulensis* populations occurring in Ubatuba which are intensively exploited by trawling.

MATERIALS AND METHODS

Study area

Located on the northern coast of São Paulo state (southeastern Brazil), Ubatuba Bay (23°26’S, 45°01’W) is a traditional seabob shrimp fishing area, with more than 100 small-scale trawling boats operating (IP/APTA/SAA/SP2021). With a total area of 8 km², the bay is 4.5 km wide at its entrance and decreases in width towards the beach (Mantelatto and Fransozo 1999). Three water masses influence the region: Coastal Water (Temperature > 20°C

and Salinity < 36); Tropical Water (Temperature > 20°C and Salinity > 36); and South Atlantic Central Water (Temperature < 18°C and Salinity < 36) (Castro-Filho et al. 1987). The seasonal influence of the water masses, coupled with the several bays formed by the presence of terminal spurs of Serra do Mar (Suguio and Martin 1978), and the transition of the tropical zone to the temperate zone, make the Ubatuba region a biodiversity hotspot for several marine organisms.

Data collection

Samples were collected monthly from January to December 2000. Shrimps were captured with a fishing boat equipped (10 m) with double rig nets, with 20 mm mesh in the wings and 18 mm mesh at the cod end, towed at an average speed of 1.7 knots. Nine transects (2 km each) were trawled: four in the internal area of the Ubatuba Bay (2, 5, 10 and 15 m isobaths) and five in the external area (20, 25, 30, 35 and 40 m

isobaths) (Figure 1). Each trawl was carried out for 30 min and an ecobathymeter coupled with a GPS was used to record the depth of sampling sites.

The captured shrimps were stored in labeled plastic bags and kept frozen until analysis. In the laboratory, they were identified according to Pérez-Farfante and Kensley (1997) and Costa et al. (2003) and sexed. Afterwards, the carapace length (CL in mm) of each individual was measured with a 0.01 mm digital caliper, which was the linear distance between the postero-orbital margin and the median notch of the posterior margin of the carapace. It was decided to use CL because structures like the rostrum and telson can be damaged during the capture of organisms and the abdomen is flexible (Cole and Mistakidis 1953). Individuals with carapace length less than 25 mm were considered juveniles (Costa et al. 2008).




Figure 1. Map of the study area indicating the location and depth of transects.

Data analysis

The growth and longevity of each species were analyzed separately for males and females. For each sampling month, the frequency of CL was distributed in 1 mm size classes. Modes were calculated using PeakFit software (PeakFit v4.12 for Windows Copyright 2000-2003 SYSTAT Software Inc.) which fits the observed frequencies to normal distribution curves so the mode and adjusted mean are the same. The “Automatic peak detection and fitting (I)” tool considering the Gaussian distribution described by the equation: $Y = a_0[\exp(-1/2(x-a_1)/a_2)^2]$, where a_0 is the amplitude, a_1 is the mean, and a_2 is the standard deviation was used to detect the central values of each mode. All the chosen cohorts were adjusted to the von Bertalanffy growth function (VBGF) $CL_t = CL_\infty(1 - \exp^{-k(t-t_0)})$, in which CL_t is the carapace length at time (i.e., age) t ; CL_∞ is the asymptotic length, k is the growth coefficient and t_0 is the theoretical age at zero length (von Bertalanffy 1938). These growth parameters were estimated for the different cohorts using Solver supplement in Microsoft Excel, varying in the equation: CL_∞ , k and t_0 , which minimizes the sum of residuals between the lengths observed in the field and those calculated by the von Bertalanffy model (Miazaki et al. 2021). Cohorts that presented a coherent biological rhythm in relation to longevity, k and CL_∞ , were chosen. An F test ($P = 0.05$) was applied to compare growth curves between sexes, and assess if a single curve could describe these species growth (Cerrato 1990). The longevity was estimated through the von Bertalanffy inverse equation modified by D’Incao and Fonseca (1999), in which $t_0 = 0$ and $CL/CL_\infty = 0.99$ and the given longevity equation is $t = (t_0 - (1/k) \ln(1 - CL_t/CL_\infty))$.

Length-converted catch curves: $-\ln(N/\Delta t) = a + bt$, were used to estimate the total mortality coefficient (Z), in which b is the regression slope that estimates Z , N is the number of individuals in each size class estimated by unit effort of capture h^{-1} , Δt is the time for the growth of individuals across this size class and t is the average age of the individual in this size class (Pauly 1990). Catch curves were estimated for each sex and compared by covariance analysis (ANCOVA) (Fonseca and D’Incao 2006). Mortality coefficient (Z) was defined as the sum of the fishing mortality coefficient (F) caused by fishing operations, and the natural mortality

coefficient (M) including predation, competition, disease, and adverse environmental conditions ($Z = F+M$).

Natural mortality (M) was estimated based on the mean value found from the methods of Pauly (1980) and Taylor (1959, 1960). According to Pauly: $\log(M) = -0.0066 - 0.279 \log(CL_\infty) + 0.6543 \log(k) + 0.4634 \log(T)$, in which CL_∞ and k are the growth parameters obtained by VBGF and T is the annual average temperature ($^{\circ}C$) of the habitat. The annual average bottom temperature was 20.91 ± 2.57 $^{\circ}C$. Taylor correlates M , k and t_0 : $M = -\ln[(1-0.95)/A_{0.95}]$, where $A_{0.95}$ is the age at which the individual reaches 95% of its asymptotic size and is defined by: $A_{0.95} = t_0 - (2.996/k)$. Fishing mortality (F) was calculated by the difference between Z and M ($F = Z - M$). The exploitation rate (E) was calculated by the ratio between F and Z : $E = F/Z$ (Sparre and Venema 1997).

RESULTS

A total of 1,231 individuals of *F. brasiliensis* and 687 individuals of *F. paulensis* were analyzed (Table 1). For *F. brasiliensis* the CL ranged from 7.50 to 35.70 mm (19.14 ± 5.66 mm) in males, and from 6.60 to 45.50 mm (18.95 ± 6.63 mm) in females (Figure 2). As for *F. paulensis*, the CL ranged from 10.40 to 35.00 mm (17.46 ± 4.26 mm) in males, and from 9.10 to 46.60 mm (18.52 ± 6.12 mm) in females.

Table 1. Abundance of *Farfantepenaeus brasiliensis* and *F. paulensis* individuals per isobath captured from January to December 2000 in Ubatuba region.

Isobath	Species	
	<i>F. brasiliensis</i>	<i>F. paulensis</i>
2 m	249	149
5 m	522	441
10 m	14	16
15 m	57	2
20 m	14	6
25 m	31	30
30 m	101	16
35 m	220	20
40 m	23	7
Total	1,231	687

Nine modal groups were selected for each sex of *F. brasiliensis* (Figure 3). The pooled

growth curves for males resulted in estimates of $CL_{\infty} = 41.08$ mm, $k = 0.198$ month⁻¹ (2.41 year⁻¹), $t_0 = -0.062$, and for females of $CL_{\infty} = 47.32$ mm, $k = 0.183$ month⁻¹ (2.23 year⁻¹), $t_0 = -0.376$ (Figure 4).

Maximum longevity (t_{max}) was estimated at 697 days (1.91 years) for males and 748 days (2.05 years) for females. Statistical comparison (F test) between estimated curves for both sexes showed significant differences ($F_{calc} = 56.10 > F_{tab} = 2.72$; $GL = 76$; $P < 0.001$).

For *F. paulensis*, six modal groups for males

and five for females were selected. The pooled growth curve of each sex resulted in estimates of $CL_{\infty} = 36.55$, $k = 0.198$ month⁻¹ (2.41 year⁻¹), $t_0 = -0.356$ for males and $CL_{\infty} = 49.24$ mm, $k = 0.207$ month⁻¹ (2.51 year⁻¹), $t_0 = -0.269$ for females (Figure 5). The t_{max} was estimated at 697 days (1.91 years) for males and 660 days (1.81 years) for females. Statistical comparison (F test) between estimated curves for both sexes showed significant differences ($F_{calc} = 304.62 > F_{tab} = 2.80$; $GL = 48$; $P < 0.001$).




Figure 2. Size frequency distribution of males and females of *Farfantepenaeus brasiliensis* and *F. paulensis* sampled monthly, from January to December 2000, in Ubatuba region.




Figure 3. Modal groups selected for males and females of *Farfantepenaeus brasiliensis* and *F. paulensis* sampled monthly, from January to December 2000, in Ubatuba region. The lines represent the modal groups.




Figure 4. Growth curves and parameters of von Bertalanffy equation estimated separately for males and females of *Farfantepenaeus brasiliensis* sampled monthly from January to December 2000 in Ubatuba region. The centerline is the mean and outer lines are prediction intervals (95%).




Figure 5. Growth curves and parameters of von Bertalanffy equation estimated separately for males and females of *Farfantepenaeus paulensis* sampled monthly from January to December 2000 in Ubatuba region. The centerline is the mean and outer lines are prediction intervals (95%).

Total mortality (Z) for *F. brasiliensis* was estimated as 10.44 year⁻¹ and 10.71 year⁻¹ for males and females, respectively, and estimated at 10.16 year⁻¹ for males and 12.77 year⁻¹ for females of *F. paulensis*. There were no statistical differences in Z between sexes of both species

and between the species (ANCOVA, *P* > 0.05) (Table 2, Figure 6). The estimated values for the natural (M) and fishing (F) mortality coefficients and exploitation rates (E) by sex and species are in Table 2. Estimates of E for both sexes and for both species were higher than 0.5 (Table 2).

Table 2. Annual estimates (year⁻¹) of natural mortality (M), fishing mortality (F), total mortality (Z), exploitation rates (E) and Standard Deviation (SD) by sex for *Farfantepenaeus brasiliensis* and *F. paulensis*.

Species	Sex	M	F	Z	E
<i>F. brasiliensis</i>	♂	2.47	7.97	10.44	0.76
	♀	2.28	8.42	10.71	0.79
	Mean ± SD	2.38 ± 0.13	8.20 ± 0.32	10.58 ± 0.19	0.78 ± 0.02
<i>F. paulensis</i>	♂	2.52	7.64	10.16	0.75
	♀	2.52	10.25	12.77	0.80
	Mean ± SD	2.52 ± 0	8.95 ± 1.85	11.47 ± 1.85	0.78 ± 0.04




Figure 6. Length-converted catch curves estimated separately for males and females of *Farfantepenaeus brasiliensis* and *F. paulensis* sampled monthly from January to December 2000 in Ubatuba region.

DISCUSSION

In terms of growth, mortality and stock exploitation, congeners *F. brasiliensis* and *F. paulensis* presented similar values for these parameters in Ubatuba Bay. Overall, females of both species reached larger asymptotic sizes than males. This sexual dimorphism is common for Penaeidae (Dall et al. 1990) and has been observed in *F. subtilis* (Santos et al. 2020), *L. schmitti* (Miazaki et al. 2018) and *Penaeus merguensis* de Man, 1888 (Saputra et al. 2018). It is related to the family reproductive strategy, i.e., pure searching. As these shrimps live aggregated in high densities, males do not guard females or territories prior to mating, so most of their energy may be allocated in developing faster in order to reproduce (Bauer 2010). As for females, they grow larger because the need to allocate their ovaries and other reproductive structures (Crisp et al. 2017).

The values in our study regarding CL_{∞} were smaller compared with previous studies with the same species. We found values of CL_{∞} of 41.08 mm for males and 47.32 mm for females of *F. brasiliensis*; and 36.55 mm for males and 49.24 mm for females of *F. paulensis*. Leite Jr and Petrere Jr (2006) found for *F. brasiliensis* values of CL_{∞} of 50.15 mm for males and 45.91 mm for females, and for *F. paulensis* values of 71.07 mm for males and 61.74 mm for females. It is important to highlight that as we sampled in a coastal bay (up to 20 m deep) and in its adjacent marine area (up to 40 m deep) on the northern coast of the São Paulo state, Leite Jr and Petrere Jr op. cit. obtained data from the fishing fleet located on the state's southern coast, which usually operates in greater depths and captures larger individuals. The estimation of L_{∞} is heavily influenced by the maximum length of the species in the sample and therefore by the sampling area (Froese and Binohlan 2000).

According to Vogt (2019), the average longevity of Penaeidae is 2.1 years. In our study, we found longevities of 1.91 years for males and 2.05 years for females of *F. brasiliensis*, and 1.91 years for males and 1.81 years for females of *F. paulensis*. While males of both species presented the same longevity, there was a considerable discrepancy for females. Females of *F. brasiliensis* may live up to three months longer

than males in general, but females of *F. paulensis* live a month less than males. This is especially concerning for the species in the area if these females are captured in their reproductive period before spawning. Close values of longevity were obtained in northeastern Brazil by Silva et al. (2015) and Santos et al. (2020) for *F. subtilis* and by Nwosu (2009) for *Farfantepenaeus notialis* (Pérez-Farfante, 1967) in Nigeria.

Growth rates (k) for *F. brasiliensis* and *F. paulensis* remained above 1.8 year^{-1} as suggested by Garcia and Le Reste (1981) for penaeids. The k of females of both species were very close to the values found by Mello (1973) in São Paulo state (23°S), from 2.52 year^{-1} for *F. brasiliensis* and 2.4 year^{-1} for *F. paulensis*. However, we found much higher k values for males than Mello op. cit, from 1.62 year^{-1} for *F. brasiliensis* and 1.25 year^{-1} for *F. paulensis*. Considerably higher k values were found herein than those found by D'Incao (1984) for both sexes of *F. paulensis* (1.03 year^{-1} for females and 1.25 year^{-1} for males) in Lagoa dos Patos - Rio Grande do Sul state (31°S), and by Leite Jr and Petrere Jr (2006) in Santos - São Paulo state (23°S) for both species, which were 0.9 year^{-1} for females and 0.84 year^{-1} for males of *F. brasiliensis* and 1.1 year^{-1} for females and 0.83 year^{-1} for males of *F. paulensis*. Values found in our study were probably higher than those found by Leite Jr and Petrere Jr op. cit because they obtained data from fishery terminals, which usually discard small individuals of little economic value. We also sampled during the closed season; therefore, more juvenile individuals were included in the analysis.

An organism's growth rate can vary due to both intrinsic (genetic effects) and extrinsic reasons, e.g., variation in food availability, temperature, oceanographic phenomena (Alford and Jackson 1993), as well as from fishing pressure. According to D'Incao (1984), results revealed by Mello (1973) were probably due to the increased fishing fleet in São Paulo state coinciding with its sampling period, thus the increase in fishing may be associated with the reduced number of individuals reaching adulthood, consequently decreasing the individual length of catches. This is also probably occurring in our study area.

Natural mortality (M) was slightly higher for males, which is probably a reflex of their smaller sizes, since smaller individuals are more likely

to be predated (Fonteles-Filho 2011). Fishery mortality (F) was higher in females. Overall, F varies with total length of individuals and is directly related to the effort and choice of fishing gear (Fonteles-Filho 2011). However, since males of both species are smaller, they present lower F values compared to females. The F values found in this study were higher than those found by Leite Jr and Petrere Jr (2006), who recorded 3.8 year⁻¹ for males and 2.8 year⁻¹ for females of *F. brasiliensis*, and 4.7 year⁻¹ for males and 3.4 year⁻¹ for females of *F. paulensis* in the Santos region, on the south-central coast of São Paulo. Nwosu (2009) studied exploited stocks of *F. notialis* and found an F of 6.44 year⁻¹. Considering that the majority of individuals sampled were juveniles (i.e. CL < 25 mm) (see Figure 2), such a high F is extremely harmful to the stock's maintenance.

Globally speaking, Penaeid fisheries are responsible for high fishing mortality rates contributing to the high Z values found for populations of economic valuable species (Garcia and Le Reste 1981). Many stocks of targeted shrimps in tropical fisheries have been over-exploited. According to Gulland (1970) and Francis (1974) the optimum exploitation rate (E) for a fishery stock is around 0.5 when $F_{opt} = M$. Studies available in the literature, such as Niamaimandi et al. (2007) on *Penaeus semisulcatus* De Haan, 1844, Safaie (2015) on *P. merguensis*, Saputra et al. (2019) on *Penaeus indicus* H. Milne Edwards, 1837, and De Carvalho et al. (2019) on *L. schmitti*, resulted in high estimates of Z and all stocks were explored over the optimum sustainable yield.

In our study, F values were considerably higher than M values. During the sampling period, E values were higher than 0.7 for both sexes of both species. Therefore, pink-shrimp stocks were over-exploited in the Ubatuba region. It is important to consider that, due to the small estuaries, *F. paulensis* finishes its juvenile growth in the bays of Ubatuba. On the other hand, *F. brasiliensis* does not enter the estuaries and completely develops in shallower portions (Costa et al. 2008). Therefore, juveniles and pre-adults who have not yet migrated to the adult stock and have not reproduced are vulnerable to the seabob fisheries, which occur at up to 20 m deep, especially if the closed season is not suitable for the species. Miazaki et al. (2021) recorded a high exploitation rate for

Xiphopenaeus spp. in the same area, reinforcing pink shrimps' vulnerability.

Seasonal closures are an important measure adopted by many tropical and subtropical shrimp fisheries to control fishing effort and prevent over-exploitation (Watson and Restrepo 1995). After 20 years of closed seasons, there still have been no recent studies about the area regarding the growth, longevity and mortality of pink shrimps. Such studies are urgently needed since several adjustments of fishing closure that occurred in the Southeast and South of Brazil (IBAMA 2008) in 2008 are still being enforced today. Data obtained from the IP/APTA/SAA /SP (2022), show that landings of pink shrimps captured in Ubatuba in the past 20 years ranged from 1.357 t in 2006 to 86 t in 2018, with annual averages of 50.308 ± 21.430 t. In 2000, 40.661 t of pink shrimp were captured in the area, while in 2021, similar values were registered, at 42.180 t. Thus, our results are of paramount importance and necessary not only to make comparisons with the current stock situation but also to reinforce the importance of regulations that control fishing activities, seeking to prevent further collapses of natural stocks, minimize the impacts caused by fishing and maintain ecologically sustainable level of stocks.

ACKNOWLEDGEMENTS

We are grateful to the staff of the Laboratório de Camarões Marinhos e de Água Doce (LABCAM) and the Núcleo de Estudos em Biologia, Ecologia e Cultivo de Crustáceos (NEBECC) for assistance in fieldwork. This research was supported by a grant from the São Paulo Research Foundation – FAPESP (#2019/01308-5, #97/12108-6, #97/12106-3, #97/12107-0) and Thematic Biota INTERCRUSTA (#2018/13685-5); by a grant from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES (#88887.161311/2017-00); and by the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (Research Fellowship PQ 306672/2018-9).

 REFERENCES

- ALFORD RA, JACKSON GD. 1993. Do cephalopods and larvae of other taxa grew asymptotically? *Am Nat.* 141 (5): 717-728. doi:10.1086/285501.
- BARIOTO JG, STANSKI G, GRABOWSKI RC, COSTA RC, CASTILHO AL. 2017. Ecological distribution of *Penaeus schmitti* (Dendrobranchiata: Penaeidae) juveniles and adults on the southern coast of São Paulo state, Brazil. *Mar Biol Res.* 13 (6): 693-703. doi:10.1080/17451000.2017.1287923.
- BAUER RT. 2010. Chemical Communication in Decapod Shrimps: The Influence of Mating and Social Systems on the Relative Importance of Olfactory and Contact Pheromones. In: BREITHAAPT T, THIEL M, editors. *Chemical Communication in Crustaceans*. Springer, New York, NY. doi:10.1007/978-0-387-77101-4_14.
- BRANCO JO, LUNARDON-BRANCO MJ, FINIS A. 1994. Crescimento de *Xiphopenaeus kroyeri* (Heller, 1862) (Crustacea: Natantia: Penaeidae) da região de Matinhos, Paraná, Brasil. *Arch Biol Technol.* 37 (1): 1-8.
- CARVALHO C, OSHIRO LMY, KEUNECKE KA. 2021. Growth and mortality analyses of the white shrimp *Penaeus schmitti* (Decapoda: Penaeidae) in Sepetiba Bay, Brazil: An exploited data-deficient species. *Reg Stud Mar Sci.* 42: 101641. doi:10.1016/j.rsma.2021.101641.
- CASTILHO AL, BAUER RT, FREIRE FAM, FRANZOZO V, COSTA RC, GRABOWSKI RC, FRANZOZO A. 2015. Lifespan and reproductive dynamics of the commercially important sea bob shrimp *Xiphopenaeus kroyeri* (Penaeoidea): synthesis of a 5-year study. *J Crust Biol.* 35 (1): 30-40. doi:10.1163/1937240X-0002300.
- CASTRO-FILHO BM, MIRANDA LB, MYAO SY. 1987. Condições hidrográficas na plataforma continental ao largo de Ubatuba: variações sazonais e em média escala. *Bolm Inst Oceanogr.* 35 (2): 135-151.
- CERRATO RM. 1990. Interpretable statistical tests for growth comparisons using parameters in the von Bertalanffy equation. *Can J Fish Aquat.* 47 (7): 1416-1426. doi:10.1139/f90-160.
- COLE HA, MISTAKIDIS MN. 1953. A device for the quick and accurate measurement of carapace length in prawns and shrimps. *ICES J Mar Sci.* 19 (1): 77-79. doi:10.1093/icesjms/19.1.77.
- COSTA RC, FRANZOZO A. 1999. A nursery ground for two tropical pink-shrimp *Penaeus* species: Ubatuba bay, northern coast of São Paulo, Brazil. *Nauplius.* 7: 73-81.
- COSTA RC, FRANZOZO A, MELO GAS, FREIRE FAM. 2003. An illustrated key for Dendrobranchiata shrimps from the northern coast of São Paulo, Brazil. *Biota Neotrop.* 3 (1): 1-12. doi:10.1590/S1676-06032003000100011.
- COSTA RC, LOPES M, CASTILHO AL, FRANZOZO A, SIMÕES SM. 2008. Abundance and distribution of juvenile pink shrimps *Farfantepenaeus* spp. in a mangrove estuary and adjacent bay on the northern shore of São Paulo State, southern Brazil. *Invertebr Reprod Dev.* 52 (1-2): 51-58. doi:10.1080/07924259.2008.9652272.
- COSTA RC, BOCHINI GL, SIMÕES SM, LOPES M, SANCINETTI G, CASTILHO AL, FRANZOZO A. 2016. Distribution pattern of juveniles of the pink shrimps *Farfantepenaeus brasiliensis* (Latreille, 1817) and *F. paulensis* (Pérez-Farfante, 1967) on the southern Brazilian coast. *Nauplius.* 24: 1-10. doi:10.1590/2358-2936e2016024.
- CRIALES MM, BROWDER JA, MOOERS CNK, ROBBLEE MB, CARDENAS H, JACKSON TL. 2007. Cross-shelf transport of pink shrimp larvae: interactions of tidal currents, larval vertical migrations and internal tides. *Mar Ecol Prog Ser.* 345: 167-184. doi:10.3354/meps06916.
- CRISP JA, FRANCES DSML, TWEEDLEY JR, PARTRIDGE GJ, MOHEIMANI NR. 2017. Quantitative determination of ovarian development in penaeid prawns (Decapoda: Penaeidae). *J Crust Biol.* 37 (1): 81-89. doi:10.1093/jcbiol/ruw001.
- DALL W, HILL BJ, ROTHILSBURG PC, STAPLES DJ. 1990. The biology of the Penaeidae. In: BLAXTER JHS, SOUTHWARD AJ, editors. *Advances in Marine Biology*. San Diego: Academic Press. 489 p.
- DE CARVALHO C, KEUNECKE KA, LAVRADO HP. 2019. Morphometric variation in pink shrimp populations at Rio de Janeiro coast (SE Brazil): are they really similar in closer areas? *An Acad Bras Cienc.* e20180252. doi:10.1590/0001-3765201920180252.

- D'INCAO F. 1984. Estudo sobre o crescimento de *Penaeus paulensis* Perez-Farfante, 1967 da Lagoa dos Patos, RS, Brasil, Decapoda, Penaeidae. *Atlântica*. 7: 73-84.
- D'INCAO F. 1991. Pesca e biologia de *Penaeus paulensis* na lagoa dos Patos, RS, Brasil. *Atlântica*. 13: 159-169.
- D'INCAO F, FONSECA DB. 1999. Performance of the von Bertalanffy growth curve in penaeid shrimp: a critical approach. Proceedings of the fourth international crustacean congress, Amsterdam, The Netherlands. 5 p.
- D'INCAO F, VALENTINI H, RODRIGUES LF. 2002. Avaliação da pesca de camarões nas regiões Sudeste e Sul do Brasil. 1965-1999. *Atlântica*. 24: 49-62.
- FONSECA DB, D'INCAO F. 2006. Mortality of *Kalliapseudes schubartii* in unvegetated soft bottoms of the estuarine region of the Lagoa dos Patos. *Arch Biol Technol*. 49 (2): 257-261.
- FONTELES-FILHO AA. 2011. Oceanografia, biologia e dinâmica populacional de recursos pesqueiros. Fortaleza: Expressão Gráfica e Editora. 460 p.
- FRANCIS RC. 1974. Relationship of Fishing Mortality to Natural Mortality at the Level of Maximum Sustainable Yield Under the Logistic Stock Production Model. *J Fish Res Board Can*. 31 (9): 1539-1543.
- FRANCO ACNP, SCHWARZ JR R, PIERRI N, SANTOS GC. 2009. Levantamento, sistematização e análise da legislação aplicada ao defeso da pesca de camarões para as regiões sudeste e sul do Brasil. *Bol Inst Pesca*. 35 (4): 687-699.
- FROESE R, BINOHLAN C. 2000. Empirical relationships to estimate asymptotic length, length at first maturity and length at maximum yield per recruit in fishes, with a simple method to evaluate length frequency data. *J Fish Biol*. 56: 758-773. doi:10.1006/jfbi.1999.1194.
- FROESE R, TSIKLIRAS CA, STERGIOU KI. 2011. Editorial note on weight-length relations of fishes. *Acta Ichthyol Piscat*. 41: 261-263.
- GARCIA S, LE RESTE L. 1981. Lyfe cycles, dynamics, exploration and management of coastal penaeid shrimp stocks. *FAO Fish Tec Pap*. 203. 215 p.
- GRABOWSKI RC, SIMÕES SM, CASTILHO AL. 2014. Population structure, sex ratio and growth of the seabob shrimp *Xiphopenaeus kroyeri* (Decapoda, Penaeidae) from coastal waters of southern Brazil. *ZooKeys*. 457: 253-269. doi:10.3897/zookeys.457.6682.
- GULLAND JA. 1970. The fish resources of the ocean. *FAO Fish. Tech. Pap*. 97. 425 p.
- GULLAND JA, ROTHSCHILD BJ. 1984. Penaeid shrimps: their biology and management. England: Fishing News Books. 308 p.
- HILBORN R, MINTE-VERA CV. 2008. Fisheries-induced changes in growth rates in marina fisheries: are they significant? *Bull Mar Sci*. 83 (1): 95-105.
- IBAMA. 2008. Instrução Normativa No 189. 23 de setembro de 2008. Pub. L. No. Process IBAMA/SC number: 2026.001828/2005-35.
- IP/APTA/SAA/SP. 2021. Estatística Pesqueira Marinha e Estuarina do Estado de São Paulo. Consulta On-line. Programa de Monitoramento da Atividade Pesqueira Marinha e Estuarina do Estado de São Paulo. Instituto de Pesca (IP), Agência Paulista de Tecnologia dos Agronegócios (APTA), Secretaria de Agricultura e Abastecimento do Estado de São Paulo (SAA/SP). [accessed: 2022 January 10]. <http://www.propesq.pesca.sp.gov.br/>.
- KILADA R, DRISCOLL JG. 2017. Age determination in crustaceans: a review. *Hydrobiologia*. 799: 21-36. doi:10.1007/s10750-017-3233-0.
- KING M. 2007. Fisheries biology, assessment and management. Oxford: Blackwell Publishing. 400 p.
- KJERFVE B. 1986. "Comparative Oceanography of Coastal Lagoons." In: Wolfe DA, editor. *Estuarine Variability*. New York: Academic Press. p. 63-81. doi:10.1016/B978-0-12-761890-6.50009-5.
- LAMPARELLI ML, COSTA MP, PRÓSPERI VA, BEVILÁQUA JE, ARAÚJO RPA, EYSINK GGL, POMPÉIA S. 2001. Sistema Estuarino de Santos e São Vicente. Relatório Técnico CETESB. São Paulo. 178 p.
- LEITE JR NO, PETRERE JR M. 2006. Growth and mortalities of the pink-shrimp *Farfantepenaeus brasiliensis* (Latreille, 1970) and *F. paulensis* (Pérez-Farfante, 1967) in southeast Brazil. *Braz J Biol*. 66 (2): 523-536.
- LOPES M. 2012. Distribuição e dinâmica populacional dos camarões-rosa, *Farfantepenaeus brasiliensis* (Latreille, 1817) e *F. paulensis* (Pérez-Farfante, 1967) e do camarão branco *Litopenaeus schmitti* (Burkenroad, 1936) (Decapoda: Dendrobranchiata: Penaeidae) no complexo baía-estuário de Santos-São Vicente, São Paulo, Brasil: Subsídios científicos para a averiguação do

- período ideal de defeso. [PhD Thesis] Botucatu: Universidade Estadual Paulista Júlio de Mesquita Filho, Instituto de Biociências. <http://hdl.handle.net/11449/106483>.
- LOPES DFC, SILVA EFB, PEIXOTO SRM, FRÉDOU FL. 2014. Population biology of the seabob-shrimp *Xiphopenaeus kroyeri* (Heller, 1862) captured on the south coast of Pernambuco state, northeast Brazil. *Braz J Oceanogr.* 62 (4): 331-340. doi:10.1590/S1679-87592014079706204.
- MANTELATTO FLM, FRANSOZO A. 1999. Characterization of the physical and chemical parameters of Ubatuba Bay, northern coast of São Paulo State, Brazil. *Rev Bras Biol.* 59: 23-31.
- MANTELATTO FLM, BERNARDO CH, SILVA TE, BERNARDES VP, COBO VJ, FRANSOZO A. 2016. Composição e distribuição de crustáceos decápodes associados à pesca do camarão-sete-barbas *Xiphopenaeus kroyeri* (Heller, 1862) no litoral norte do estado de São Paulo. *Bol. Inst. Pesca.* 42: 307-326. doi.org/10.20950/1678-2305.2016v42n2p307.
- MELLO JTC. 1973. Estudo populacional do camarão-rosa, *Penaeus brasiliensis* Latreille, 1817, e *Penaeus paulensis* Pérez-Farfante, 1967. *Bol Inst Pesca.* 2 (2): 19-65.
- MIAZAKI LF, SANTOS APF, SALVATI DS, SIMÕES SM, COSTA RC. 2018. Growth, longevity, and juvenile recruitment of the white shrimp *Litopenaeus schmitti* (Burkenroad, 1936) (Decapoda: Penaeoidea) in southeastern Brazil. *Acta Sci Biol Sci.* 40 (1): e40379. doi: 10.4025/actascibiolsoci.v40i1.40379.
- MIAZAKI LF, HECKLER GS, SANTOS APF, CASTILHO AL, PESPINELLI RA, COSTA RC. 2021. Growth parameters, longevity, and mortality of the seabob shrimp *Xiphopenaeus* spp. (Decapoda: Penaeidae) in four important fishing regions of the southeastern Brazil. *Fish Oceanog.* 00: 1-16. doi:10.1111/fog.12533.
- MUSIELLO-FERNANDES J, ZAPPES CA, HOSTIM-SILVA M. 2017. Small-scale shrimp fisheries on the Brazilian coast: Stakeholders perceptions of the closed season and integrated management. *Ocean Coast Manage.* 148: 89-96. doi.org/10.1016/j.ocecoaman.2017.07.018.
- NIAMAIMANDI N, ARSHAD AB, DAUDB SK, SAED RC, KIABI B. 2007. Population dynamic of green tiger prawn, *Penaeus semisulcatus* (De Haan) in Bushehr coastal waters, Persian Gulf. *Fish Res.* 86: 105-112. doi:10.1016/j.fishres.2007.05.007.
- NWOSU FM. 2009. Population Dynamics of the Exploited Penaeid Shrimp, *Penaeus (Farfantepenaeus) notialis* in the Cross River Estuary, Nigeria. *J Fish Int.* 4 (4): 62-67. doi: 10.3923/jfish.2009.62.67.
- PAULY D. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *ICES J Mar Sci.* 39 (2): 175-192. doi:10.1093/icesjms/39.2.175.
- PAULY D. 1990. Length-converted catch curves and the seasonal growth of fishes. *Fishbyte.* 8 (3): 24-29.
- PÉREZ-FARFANTE I, KENSLEY B. 1997. Penaeoid and Sergestoid Shrimps and Prawns of the World. Keys and diagnoses for the families and genera. Éditions du Muséum National d'Histoire Naturelle. Paris. 233 p.
- IP/APTA/SAA/SP. 2022. Estatística Pesqueira Marinha e Estuarina do Estado de São Paulo. Consulta On-line. Programa de Monitoramento da Atividade Pesqueira Marinha e Estuarina do Estado de São Paulo. Instituto de Pesca (IP), Agência Paulista de Tecnologia dos Agronegócios (APTA), Secretaria de Agricultura e Abastecimento do Estado de São Paulo (SAA/SP). [accessed: 2022 January 20]. <http://www.propesq.pesca.sp.gov.br/>.
- REIS JR JJDC, FREIRE KMF, ROSA LC, BARRETO TMRDR, PAULY D. 2019. Population dynamics of Atlantic seabob *Xiphopenaeus kroyeri* (Decapoda: Penaeidae) off the state of Sergipe, north-eastern Brazil. *J Mar Biol Assoc UK.* 99 (1): 143-153. doi:10.1017/S0025315417001916.
- SANTOS RC, PERROCA JF, COSTA RC, HIROSE GL. 2020. Population dynamics of *Farfantepenaeus subtilis* (Pérez-Farfante, 1967) and *Litopenaeus schmitti* (Burkenroad, 1936) (Decapoda: Penaeidae) and evidence of habitat partitioning in the northeast of Brazil. *Reg Stud Mar Sci.* 35: 101218. doi: 10.1016/j.rsma.2020.101218.
- SAFAIE M. 2015. Population dynamics for banana prawns, *Penaeus merguensis* de Man, 1888 in coastal waters off the northern part of the Persian Gulf, Iran. *Trop Zool.* 28 (1): 9-22. doi:10.1080/03946975.2015.1006459.
- SAPUTRA SW, SHOLICHIN A, TAUFANI WT. 2018. Growth, mortality and exploitation rate of

- Penaeus merguensis* in the North Coast of Central Java, Indonesia. Ilmu Kelautan. 23 (4): 207-214. doi: 10.14710/ik.ijms.23.4.207-214.
- SAPUTRA SW, SOLICHIN A, TAUFANI WT, RUDIYANTI S, WIDYORINI N. 2019. Growth parameter, mortality, recruitment pattern, and exploitation rate of white shrimp *Penaeus indicus* in northern coastal waters of Western Central Java, Indonesia. Biodiversitas. 20 (5): 1318-1324. doi: 10.13057/biodiv/d200511.
- SEMENSATO XEG, DI BENEDITTO APM. 2008. Population dynamic and reproduction of *Artemesia longinaris* (Decapoda, Penaeidae) in Rio de Janeiro state, south-eastern Brazil. Bol Inst Pesca. 34: 89-98.
- SILVA EF, CALAZANS N, NOLÉ L, VIANA A, SOARES R, PEIXOTO S, FRÉDOU FL. 2015. Population Dynamics of the pink shrimp *Farfantepenaeus subtilis* in Northeastern Brazil. J Crust Biol. 35 (2): 132-139. doi:10.1163/1937240X-00002325.
- SILVA SLR, SANTOS RC, COSTA RC, HIROSE GL. 2018. Growth and population structure of the seabob shrimp *Xiphopenaeus kroyeri* (Decapoda: Penaeidae) on the continental shelf of Sergipe, Brazil. J Mar Biol Assoc U K. 99 (1): 81-92. doi:10.1017/S0025315417002041.
- SOUZA FL, KJERFVE B, KNOPPERS B, LANDIM DE SOUZA WF, DAMASCENO RN. 2003. Nutrient budgets and trophic state in a hypersaline coastal lagoon: Lagoa de Araruama, Brazil. Estuar. Coast Shelf Sci. 57 (5-6): 843-858. doi:10.1016/S0272-7714(02)00415-8.
- SPARRE P, VENEMA SC. 1997. Estimação das taxas de mortalidade. In: Introdução à avaliação de mananciais de peixes tropicais, Parte 1: Manual. FAO Tech Doc sobre as pescas No. 306/1, Rev. 2. Rome, FAO. p. 119-161.
- STAPLES DJ, VANCE DJ. 1985. Short-term and longterm influences on the immigration of postlarval banana prawns *Penaeus merguensis*, into a mangrove estuary of the Gulf of Carpentaria, Australia. Mar Ecol Prog Ser. 23: 15-29.
- SUGUIO K, MARTIN L. 1978. Formações quaternárias marinhas do litoral paulista e fluminense. Paper present at the International Symposium Coastal evolution in the Quaternary, São Paulo, SP. [accessed December 2021] <https://core.ac.uk/download/pdf/39879155.pdf>.
- TAYLOR CC. 1959. Temperature and growth – the Pacific razor clam. ICES J Mar Sci. 25 (1): 93-101.
- TAYLOR CC. 1960. Temperature, growth and mortality – the Pacific Cockle. ICES. Mar Sci. 26 (1): 117-124.
- TEODORO SSA, TEROSSI M, MANTELATTO FLM, COSTA RC. 2016. Discordance in the identification of juvenile pink shrimp (*Farfantepenaeus brasiliensis* and *F. paulensis*: Family Penaeidae): An integrative approach using morphology, morphometry and barcoding. Fish Res. 183: 244-253. doi: 10.1016/j.fishres.2016.06.009.
- VALENTINI H, D'INCAO F, RODRIGUES LF, REBELO NETO JE, RAHN E. 1991. Análise da pesca do camarão-rosa (*Penaeus brasiliensis* e *Penaeus paulensis*) nas regiões sudeste e sul do Brasil. Atlântica. 13: 143-157.
- VILELA MJ, COSTA PAS, VALENTINI JL. 1997. Crescimento e mortalidade de juvenis do camarão-rosa, *Penaeus brasiliensis* Latreille, 1817, na Lagoa de Araruama, Rio de Janeiro. Rev Bras Biol. 57: 487-499.
- VOGT G. 2012. Ageing and longevity in the Decapoda (Crustacea): A review. Zool Anz. 251 (1): 1-25. doi:10.1016/j.jcz.2011.05.003
- VOGT G. 2019. A compilation of longevity data in decapod crustaceans. Nauplius. 27: e2019011. doi:10.1590/2358-2936e2019011.
- VON BERTALANFFY L. 1938. A quantitative theory of organic growth. Hum Biol. 10: 181-213.
- WATSON RA, RESTREPO VR. 1995. Evaluating closed season options with simulation for a tropical shrimp fishery. ICES Mar Sci Symp. 199: 391-398.
- ZAR JH. 2010. Biostatistical analysis. New Jersey: Prentice Hall. 718 p.