NOTE

Size structure of Areolate grouper (*Epinephelus areolatus*) from the Saudi coast of the Arabian Gulf

JESUSITO A. VICENTE*

Pangasinan State University, Binmaley Campus, Binmaley 2417, Pangasinan, Philippines



ABSTRACT. Areolate grouper, Epinephelus areolatus of the reef-associated fish species which is highly sought-after in the seafood trade. Con ently. gh market demand resulted to overexploitation and population decline of the specie his paper aimed to determine the wild. the size structure of E. areolatus from the Arabi 55 specimens of E. areolatus collected over the 12-month sampling per reve roportions of females observed throughout the year and in size class. Man **(29** m, 358.44 g) were bigger and heavier than females (28.8 cm, 326.66 g). The 'b' v from 6 to 2.88 indicated negative allometric growth. The relationship between le and weigh ved significant positive correlations with p < 0.0000 and r² values ranging be 05-97.12%.

Key words: Arabian Gulf reolate grouper, length-weight relationship.

Estructura de del medicado (Epinephelus areolatus) de la costa saudí del Golfo Arábigo

RESV. N. El co areolado, *Epinephelus areolatus*, es una de las especies de peces asociadas a los recificações y buscada en el comercio de productos del mar. Consecuentemente, una alta den el la comerció de productos del mar. Consecuentemente, una alta den el la comerció de la sobreexplotación y la disminución de la población de la especie en la nuevaleza. Este trabajo tuvo como objetivo determinar la estructura de tamaño de *E. areolatus* del Golfo objeto. Un total de 355 muestras de *E. areolatus* recolectadas durante 12 meses de muestra revelaron altas proporciones de hembras durante todo el año y en la clase de talla. Se observa que es machos (29,3 cm, 358,44 g) fueron más grandes y pesados que las hembras (28,8 cm, 326,66 g). Los valores de "b" que oscilan entre 2,86 y 2,88 indicaron un crecimiento alométrico negativo. La relación entre longitud y peso mostró correlaciones positivas significativas con p < 0,0000 y valores de r² que oscilaron entre 96,05-97,12%.

Palabras clave: Golfo Arábigo, mero areolado, relación largo-peso.

*Con tence: jesusit nte@ 'l.cor

ceived February 22

nte* 2683-7595 (print)

https://ojs.inidep.edu.ar

Journal of the Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP)



7951 (online)

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License Areolate groupers, *Epinephelus areolatus* (Forsskål, 1775), are reef-associated fishes found mainly on depths ranging from 6-200 m (Froese and Pauly 2013). They are usually inhabitants of seagrass beds or fine sediment bottoms near rocky reefs, dead coral, or alcyonarians, in shallow continental shelf waters (Tupper and Sheriff 2008; Froese and Pauly 2013; Sanaye 2014). Despite the wide distribution of Family Epinephilidae, this species is found only in the Indo-Pacific region (Russell and Houston, 1989; Heemstra and Randall 1993; Ottolenghi et al. 2004; Sanaye 2014) (Figure 1). Areolate

groupers are important fisheries and aquaculture species. Together with other grouper species, they are considered most highly sought-after food fishes in the seafood trade (Kuo 1995; Sadovy et al. 2003). However, the high demand of grouper in the market generated a negative ecological impact. It is the most

intensively exploited group in the live fish trade, implying that this group is heavily overfished (Morris et al. 2000). Studies reveal that trade often follows a pattern of sequential overexploitation where the most sought species is targeted and fished out first before the less valuable species (Johannes and Riepen 1995; Sluka 1997). Barrowman and Myers (1996) and Reinert et al. (2005) indicated that the removal of considerable number of sexually mature fish in the stock would compromise the overall stock reproductive output, especially the species forming concentrated and brief spawning aggregations (Sadovy and Domeier 2005), such as this group. Population decline was already observed in some areas ap attributed to overfishing, habitat degradation pollution, destructive fishing techniques, his

export demand, etc. (Johannes 1997; Sadovy 2000). Together with the compounding effect brought about by natural and anthropogenic causes, stocks cannot sustain their population. As a result, natural stocks are depleted or even worse, some species are threatened to become extinct. At present, some species of groupers are already considered 'threatened organisms' according to IUCN categories and criteria.

In the region, fisheries are considered as the most important natural resource next to oil (Carpenter et al. 1997). Sauc Arabia, production figures reveal an increasing and from 1996 to 2005 (Ministry of Agrey' are Saudi Arabia, 2007). Only 2005, a total of 74,779 t of seafood products here oduced from traditional fishing, industry fishing is sh and marine aquaculture. Nearly the refourths (70.5%) of total catch condited from traditional fishing. Industrial fishing and aquaculture account only for 10.2% and 19.2% respectively. There are two main grouper commonly sold in the market, namely *E. coiodes* and *E. areolatus*. Their prices range from eSD 15-25 and USD 8-11 kg⁻¹, respectively.




Figure 1. Geographical distribution of Epinephelus areolatus.

The purpose of this study was to determine the size structure of areolate grouper, *E. areolatus* from the Arabian Gulf. It particularly focuses on the length-weight relationship which can be a useful tool in management and conservation of the species in area.

Specimens of *E. areolatus* were collected on a monthly basis from Jubail Fish Market and Auction Center, a major fish landing site on the eastern province of Saudi Arabia, western part of the Arabian Gulf. A total of 355 samples were collected over the 12-month sampling period from February 2014 to January 2015. The relationship between total length (TL) and whole-body weight (BW) of *E. areolatus* was analyzed by measuring length and weight of the specimens. The statistical relationship between these parameters of fishes was determined by using the parabolic equation formulated by Froese (2006) as follows:

$$W = aL^b$$

where, W = weight of fish (g), L = lengt of (mm), a = constant and b = an expressing relationship between 1st th-wight. Such relationship when convered it to the logarithmic form gives a stright of relationship graphically. The same relation meaning oned above is written in logarithmic form as:

$$Ln W = Ln a + Lr$$

where, k and the confidence of determination r² were expand the option of the confidence limit. Statistical analyst were computed using EXEL STAT for Windows XLSTAT 2007).

The length of the gathered samples ranged between 17.1-47.1 cm, with the mean length of 28.95 cm. The weight ranged between 58.10-1,343.43 g, with the mean of 334.80 cm. It was further observed that males (29.3 cm, 358.44 g) were bigger and heavier than females (28.80 cm, 326.66 g). Maximum length and weight reported in other areas were as follows: 50.5 cm and 1.94

kg in Egypt (Abd-Allah et al. 2015), 29.5 cm in the Philippines (Gumanao et al. 2016), 49.5 cm and 1.5 kg in India (Nair et al. 2021), and 30.2 cm and 0.38 kg in Indonesia (Fadli et al. 2022).

The 'b' value was 2.88 for males while for females it was 2.86, and for combined sexes it was 2.87. Results indicated that the weight of E. areolatus increased with the increasing length (Figure 2). Analysis of regression shows that there was a significant relationship between the two variables with 0.0000 and r^2 values ranging between 96 5-97. %. Typically, growth in fish is explained by von ertallanfy curve (Hopkins 1992 auly 1 4: bling 2002) represented by an empedic signoid curve in many species 9). Leteef fishes like E. areolatus, (Ri er mibit during pelagic and juvenile growth stage however, it slows down during transitions into a hood to apportion more energy for breeding (Jobling 1994; Hutchings 2003; Claro Garcia-Arteaga 2014). In the present study, imputed 'b' values were 2.87 (combined sexes), 2.88 (males) and 2.86 (females). This is an indication that the fish shows a negative allometric growth implying that the parts of the fish grow slower compared to its body as a whole. This growth performance index is similar to the studies conducted from the Gulf of Suez (2.83), north coast of Aceh, Indonesia (2.86-3.31), southwest coast of India (2.95-3.2) and Davao Gulf, Philippines (3.03) (Abd-Allah et al. 2015; Gumanao et al. 2016; Nair et al. 2021; Fadli et al. 2022).

The relationship between length and weight showed a positive correlation, suggesting that the weight of *E. areolatus* increased with the increasing length. According to Jennings and Polunin (1997), this relationship is a morphometric measurement of how a species allocates mass allometrically. The association between the two variables can be useful to estimate standing crop biomass (Abd-Allah et al. 2015), assess species fitness overtime (Bolger and Connolly 1989) and provides information on production capabilities (Jobling 2002).




Figure 2. Length-weight relationship of *Epinephelus areolatus* from the Arabian Gulf, n = 355 ($n_{male} = 91$, $n_{female} = 264$). A) Combined sexes. B) Male. C) Female.

Similar to the reproductive capability of a species, growth is largely associated with the prevailing local environmental circumstances (Roff 2000). Many studies have uncovered that the pronounced growth variability within and between populations is due to the combination of both genetic and environmental factors such as temperature, photoperiod, pH, salinity, and food availability (Werner and Gilliam 1984; Manooch 1987; Conover 1990; Sadovy et al. 1992; Sale 1998; Lombardi-Carlson et al. 2008; Munday et al. 2008; Claro and Garcia-Arteaga 2014).

REFERENCES

- ABD-ALLAH E, EL-GANAINY A, OSMAN A. 2015. Age and growth of the areolate grouper *nephelus areolatus* from the Gulf of Suez. J Life Sci. 3: 7-12.
- BARROWMAN NJ, MYERS RA. 1996. Is so perula ment related to spawner abundance. Fig. 8 Rull, 94: 707-724.
- BOLGER T, CONNOLLY PL. 16 The section of suitable indices for the measurement and analysis of fish conditions. Fish Biol. 4: 171-182.
- CLARO R, GARCIA-A CEAS L. 2014. Growth patterns of fishes of L. Cuba shelf. In: CLARO R, LINDEM A C., PA. L. LR, editors. Ecology of the many coshes of Cuba. Smithsonian Institution. Ashington. p. 149-166.
- CONOVER DO. 17 The relation between capacity for growth and length of growing season: evidence for and implications of counter gradient variation. Trans Am Fish Soc. 119: 416-430.
- FADLI N, DAMORA A, MUCHLISIN ZA, DEWIYANTI I, RAMADHANIATY M, RAZY NM, MACUSI ED, SITI-AZIZAH MN. 2022. Length-weight relationships and condition factors of three *Epinephelus* grouper (Epinephelidae) harvested in the Northern Coast of Aceh, Indonesia. Int J Des Nat Ecodynamics. 17: 119-124.

- FROESE R. 2006. Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. J Appl Ichthyol. 22: 241-253.
- FROESE R, PAULY D. 2013. FishBase. http://www.fishbase.org.
- GUMANAO GS, SACEDA-CARDOZA MM, MUELLER B, Bos AR. 2016. Length-weight and length-length relationships of 9 Indo-pacific fish species (Teleosta from Lyao Gulf, Philippines. J Applichth). 32 277-285.
- HEEMSTRA & RANGLL J. 993. Groupers of the world fan a Serra dae, subfamily Epinephelic 1: An an acted and illustrated catalogue at the soupers, rockcod, hind, coral grouper and lyre 1 species known to date. Rome: FAO.
- HOPKI KD. 1992. Reporting fish growth: a ew of the basics 1. J World Aquac Soc. 23: 173-179.
- MUTCHINGS J. 2003. Life histories of fish. In: HART P, REYNOLDS D, editors. Handbook of fish biology and fisheries. Vol. 1. Oxford: Blackwell Science. p. 149-174.
- Jennings S, Polunin NVC. 1997. Impacts of predator depletion by fishing on the biomass and diversity of non-target reef fish communities. Coral Reefs. 16: 71-82.
- JOBLING M. 2002. Environmental factors and rates of development and growth. In: HART P, REYNOLDS D, editors. Handbook of fish biology and fisheries. Oxford: Blackwell Science. p. 97-102.
- JOBLING M, editor. 1994. Production and growth. Fish bioenergetics. London: Chapman and Hall. p. 147-150.
- JOHANNES RE. 1997. Grouper spawning aggregations need protection. SPC Live Reef Fish Information Bulletin. 3: 13-14.
- JOHANNES RE, RIEPEN M. 1995. Environmental, economic, and social implications of the live reef fish trade in Asia and the Western Pacific. Honolulu: The Nature Conservancy,
- Kuo C. 1995. The groupers. In: NASH C,

- NOVOTNY A, editors. Production of aquatic animals: fishes. Amsterdam: Elsevier. p. 305-317.
- LOMBARDI-CARLSON L, FITZHUGH G, PALMER C, GARDNER C, FARSKY R, ORTIZ M. 2008. Regional size, age and growth differences of red grouper (*Epinephelus morio*) along the west coast of Florida. Fish Res, 91: 239-251.
- MANOOCH CS. 1987. Age and growth of snappers and groupers. In: POLOVINA J, RALSTON S, editors. Tropical snappers and groupers: biology and fisheries management. Boulder: Westview Press.
- MORRIS AV, ROBERTS CM, HAWKINS JP. 2000. The threatened status of groupers (Epinephelinae). Biodivers Conserv. 9: 919-942.
- MUNDAY PL, JONES GP, PRATCHETT MS, WILLIAMS AJ. 2008. Climate change and the future for coral reef fishes. Fish Fish. 9: 261-285.
- NAIR RJ, SEETHA PK, SUNIL KTS, RADHAKRIS-HNAN M. 2021. Length weight relations per demersal reef fishes from southwest coast India. J Mar Biol Assoc India. 62-40-
- OTTOLENGHI F, SILVESTRI C, GIOTANO P, DAN-TELLI A, NEW M. 2004. Captured-wed Aquaculture. The fattening of tels, groupe tunas and yellowtails. Rome FAC 308 p.
- PAULY D. 1994. Quantita analysis of published data on the powth, analysis of published data on the powth, analysis of published data on the powth, analysis of the redbellied piranha, arasalmus nattereri (Characidae). Environ Bit Tishes. 41: 423-437.
- REINERT TR, JENNINGS CA, WILL TA, WALLIN JE. 2005. Decline and potential recovery of striped bass in a southeastern US estuary. Fisheries. 30: 18-25.
- RICKER WE. 1979. Growth rates and models. In: HOAR WS, RANDALL DJ, BRETT JR, editors. Fish physiology. Vol. 8. New York: Academic Press. p. 677-743.
- ROFF DA. 2000. Trade-offs between growth and reproduction: an analysis of the quantitative

- genetic evidence. J Evol Biol. 13: 434-445. RUSSELL BC, HOUSTON W. 1989. Offshore fishes of the Arafura Sea. Beagle. 6: 69-84.
- Sadovy Y. 2000. Regional survey for fry/fingerling supply and current practices for grouper mariculture: evaluating current status and long-term prospects for grouper mariculture in Southeast Asia. Final report to the collaborative APEC Grouper Preserve and Development Network (FWG 199).
- SADOVY Y, DOMEIF M. 20 A aggregation-fisheries sust table Reef sh fisheries as a case study Solve eefs 4: 254-262.
- SADOVY YOU GUEROL WAR ROMAN A. 1992. Age and glowth of red hind, *Epinephelus guttatus*. Prop. Rico and St. Thomas. Fish Bull. 90: 16-528.
- SALVY Y DONALDSON TJ, GRAHAM TR, M. S. AY F, MULDOON GJ, PHILLIPS MJ, PIMMER MA, SMITH A, YEETING B. 2003. While stocks last: the live reef food fish trade. Manila: Asian Development Bank. 169 p.
- SALE PF. 1998. Appropriate spatial scales for studies of reef-fish ecology. Austral J Ecol. 23: 202-208.
- SANAYE S. 2014. Reef fishes of India: India biodiversity portal species data. http://indiareef fishes.lifedesks.org/.
- SLUKA RD. 1997. The biology and ecology of grouper in Laamu Atoll, Republic of Maldives. Male: Oceanographic Society of Maldives.
- TUPPER M, SHERIFF N. 2008. Capture-based aquaculture of groupers. In: LOVATELLI A, HOLTHUS PF, editors. Capture-based aquaculture: global overview. FAO Fish Tech Pap. 508. p. 217-253.
- [XLSTAT]. 2007. Statistical Software for Excel. New York: Addinsoft. http://www.xlstat.com.
- WERNER EE, GILLIAM JF. 1984. The ontogenetic niche and species interactions in size-structured populations. Annu Rev Ecol Sys. 15: 393-425.